

MUTAGENICITY TEST ON
Combined Ethanol Extracts of Five Mammary Implant Components

IN AN IN VITRO CYTOGENETIC ASSAY
MEASURING CHROMOSOMAL ABERRATION FREQUENCIES IN
CHINESE HAMSTER OVARY (CHO) CELLS

REVISED FINAL REPORT

XII. CONCLUSION:

The combined ethanol extracts of [REDACTED] Elastomer Shell, [REDACTED]
Elastomer Shell, [REDACTED] Elastomer Shell, Leaf Valve Assembly, and
[REDACTED] Gel were considered negative for inducing chromosomal aberrations in Chinese hamster ovary cells under both the metabolic activation and nonactivation conditions of this assay.

XIII. REFERENCES:

- Evans, R.J.: Chromosomal aberrations produced by ionizing radiation. International Review of Cytology, 13:221-321, 1962.
- Goto, K., Maeda, S., Kano, Y., and Sugiyama, T.: Factors involved in differential Giemsa-staining of sister chromatids. Chromosoma, 66:351-359, 1978.
- Petry, P. and Wolff, S.: New Giemsa method for the differential staining of sister chromatids. Naturw., 251:156-158, 1974.
- Sokal, R.R., and Rohlf, F.J.: Biometry, Ed. 2. W.H. Freeman and Company, New York, 1981.
- Terasima, T. and Tolmach, L.J.: Changes in X-ray sensitivity of HeLa cells during the division cycle. Nature, 190:1210-1211, 1961.